Environmental Modelling & Software 102 (2018) 292–305

Contents lists available at ScienceDirect

Environmental Modelling & Software

journal homepage: www.elsevier.com/locate/envsoft

Perennial-GHG: A new generic allometric model to estimate biomass accumulation and greenhouse gas emissions in perennial food and bioenergy crops

A. Ledo [a, *], R. Heathcote [b], A. Hastings [a], P. Smith [a], J. Hillier [a]

[a] University of Aberdeen, United Kingdom
[b] Cool Farm Alliance, The Stable Yard, Vicarage Road, Stony Stratford, MK11 1BN, United Kingdom

ARTICLE INFO

Article history:
Received 4 October 2017
Accepted 10 December 2017

Keywords:
Above ground biomass
Below ground biomass
Carbon
Carbon dioxide
Decomposition
Greenhouse gas emissions
Modelling

ABSTRACT

Agriculture, and its impact on land, contributes almost a third of total human emissions of greenhouse gases (GHG). At the same time, it is the only sector which has significant potential for negative emissions through offsetting *via* the supply of feedstock for energy and sequestration in biomass and soils. Perennial crops represent 30% of the global cropland area. However, the positive effect of biomass storage on net GHG emissions has largely been ignored. Reasons for this include the inconsistency in methods of accounting for biomass in perennials. In this study, we present a generic model to calculate the carbon balance and GHG emissions from perennial crops, covering both bioenergy and food crops. The model can be parametrized for any given crop if the necessary empirical data exists. We illustrate the model for four perennial crops – apple, coffee, sugarcane, and *Miscanthus*– to demonstrate the importance of biomass in overall farm GHG emissions.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

Agriculture is an essential human activity but at the same time a substantial emitter of greenhouse gas (GHG) emissions (Robertson et al., 2000). With a rising global population, the need for agriculture to provide a secure food and energy supply is one of the main human challenges (Smith et al., 2010a). Agriculture contributes about 4.6–5.4 Gt CO_2-equivalent per year, which is 9–11% of global GHG anthropogenic emissions in 2010 (Tubiello et al., 2013; Smith et al., 2014), and the value approaches a third of total emissions if the indirect impacts of land use change, and land degradation (Wollenberg et al., 2013) are considered. At the same time it, and the other land based sectors, are the only ones which have significant potential for negative emissions through the sequestration of carbon and offsetting *via* the supply of feedstock for energy production.

In addition to land use change, major sources of GHG emissions from crop production include N_2O emission from the production and use the use of fertilizers (Robertson et al., 2000), methane emissions from paddy rice production and livestock (Yan et al., 2005), and the loss of stored biomass and soil carbon, all of which may in part be attributed to management. These emissions can be reduced or reversed, so management is a potential tool for GHG mitigation (Smith et al., 2008, 2014). To enable judicious management to be prescribed, sources of GHG emission first need to be identified and quantified.

Perennial crops such as fruit trees or bioenergy grasses like *Miscanthus* are often not differentiated from annual crops when estimating agricultural GHG emissions. However, in contrast to annual cropping systems which most often have positive GHG emissions, perennials may have net zero or even negative emissions (Glover et al., 2010; Robertson et al., 2000, 2016; McCalmont et al., 2015). Perennial agricultural management also reduces soil disturbance since annual cultivation is not required, and it adds more carbon inputs to the soil and improves soil conditions (Paustian et al., 2000; Cox et al., 2006). This, in turn, allows soil carbon to be stabilised, hence reducing emissions of carbon dioxide to the atmosphere *via* mineralization in those cases in which the soil is not saturated with carbon (Dawson and Smith, 2007). Besides, some perennial crops, and in particular perennial grasses like Miscanthus, are more effective at intercepting and utilizing water

* Corresponding author. University of Aberdeen, Institute of Environmental and Biological Sciences, School of Biological Sciences, St Machar Drive 23, Room G43, Aberdeen AB24 3UU, United Kingdom.
E-mail address: alicialedo@gmail.com (A. Ledo).

https://doi.org/10.1016/j.envsoft.2017.12.005
1364-8152/© 2018 Elsevier Ltd. All rights reserved.

and CO_2 resources (Dohleman and Long, 2009), and some need less or no fertilizer application (Hastings et al., 2009, 2017; Davis et al., 2012). This may have vital implications for GHG and mitigation options in the future; hence it is timely to develop generic, consistent, and scalable models to account for often overlooked biomass accumulation, particularly in perennial production systems.

Perennial crops accumulate carbon during their lifetime, in above and below ground components, and enhance organic soil carbon increase *via* root senescence and litter inputs. However, inconsistency in accounting for this stored biomass undermines efforts to assess the benefits of such cropping systems when applied at scale. Common product foot-printing standards e.g. the Publicly Available Standard 2020:2011 (PAS 2050), the EU renewable Fuel Directive (RED), and the GHG protocol for product life cycle accounting, for various reasons, do not consider soil carbon stock changes or biomass accumulation in carbon footprint calculations (Whitaker et al., 2010). The major concerns appear to be, firstly, the lack of reliable methods to quantify carbon stocks in the various plant components, and secondly, issues around permanence of the biomass carbon stored (Brandão et al., 2013). A consequence of this exclusion is that efforts to manage this important carbon stock are neglected. Detailed information on carbon balance is crucial to identify the main processes responsible for greenhouse gas emissions in order to develop strategic mitigation programmes. Perennial cropping systems represent 30% of the area of total global crop systems (Glover et al., 2010). Furthermore, they have a major role both in the global food (i.e. oil palm, coffee, fruit and cocoa) and bioenergy (i.e. *Miscanthus*, switchgrass, sugarcane, short rotation coppice) industries. At the same time, an increase in perennial crops or 'perennialization', is one of FAO's (Food and Agriculture Organization of the United Nations) strategies to enhance food security and ecosystem service delivery (Glover et al., 2010; Rai et al., 2011).

In this paper, we present a generic model, Perennial-GHG, to calculate the carbon balance and GHG emissions from perennial crops at farm level that does not require the level of site information necessary to run a detailed, process-based model. This model covers the cultivation period and the residue management for both food and bioenergy crops, also considering intercropping, the combination of two or more perennial crops. GHG emissions can be either positive (emissions to the atmosphere) or negative (carbon uptake from the atmosphere). Plant biomass is formed *via* carbon uptake from the atmosphere; consequently, it is stored as a negative GHG emission in the model while it is living material in the plant. Once the plant or plant part is removed or naturally released, it becomes a residue (see Fig. 1).

We then use this model to illustrate the importance of biomass in the estimation of overall GHG emissions from four important perennial crops - coffee, apple, *Miscanthus* and sugarcane – which were chosen to give examples from tropical and temperate regions, trees and grasses, and energy and food supply. We propose a model that has wide applicability and can be used both in research environments and for decision support among industry, farming, and NGO stakeholders, to evaluate actual agriculture practises, and support efforts to reduce the GHG intensity of agricultural products by accounting for biomass storage and decomposition, and persistence of carbon in the system. Plant biomass is in large part carbon fixed from the atmosphere by photosynthesis and stored in the plant. The model runs using inputs supplied by the farmer or land manager, including the cultivated area, crop or crops, and the main management options (the list of inputs is presented in Supplementary information S3). Importantly, yield is also an input in the Perennial-GHG model. The Perennial-GHG model does not aim to predict yield, as physiological crops and process-based models do, but to estimate biomass and GHG emissions in perennial crops based on expected/previously recorded/estimated yield.

The Perennial-GHG model is data-driven and based on allometric relationships of biomass increment as a function of time. Although physiological crop process-based models are common in agricultural research (Priesack and Gayler, 2009), the input data required, such as daily meteorological data, and internal parameters such as photosynthesis and evapotranspiration rate, means that they are not easy to apply outside the research community. Process based models can give accurate simulations of daily plant growth and yield, making them more accurate, but also more complex and computationally demanding, which makes them unsuitable for use by farmers/land-managers, and unsuitable for inclusion in most decision support systems.

Contrary to natural ecosystems, the shape of the trees in farmland is mainly the result of the management actions, i.e. pruning, and controlled by climatic conditions to a lesser extent. At the end of the crop cycle, tree woody biomass often reflects human actions. The generic model we are presenting is composed of two simple sub-models, to cover grasses and other perennial plants. The first is a generic individual-based sub-model (IBM) covering both woody crops in which the yield is the fruit and the plant biomass is an unharvested residue, and short rotation coppice (SRC). Trees, shrubs and climbers fall into this category. The second model is a generic area-based sub-model (ABM) covering perennial grasses, in which the harvested part includes some of the plant parts in which the carbon storage is accounted. Most second generation perennial bioenergy crops fall into this category. Both generic sub-models presented in this paper can be parametrized for different crops, and we have parametrized the sub-models for a list of crops using published empirical data. The model can also account for different varieties, geographical locations and rate of applied fertilizer, and for fine-scale analysis, it can be parametrized at farm level.

For use outside the research community, so-called "carbon calculators" have been developed. Although there are several of these, the accounting for stored biomass is relatively limited (Whittaker et al., 2013). The models we develop in this study have been co-designed with the Cool Farm Alliance to be ready for insertion in to the Cool Farm Tool (CFT, www.coolfarmtool.org) - a free-to-use, farmer-oriented GHG calculator, which has been widely used globally by industry and farming to assess GHG emissions, and identify positive interventions to mitigate GHG emissions. The CFT performed best among all farm GHG emissions calculators in the UK (Whittaker et al., 2013), and the incorporation of improved accounting for biomass in perennials will enable wider use in the bioenergy sector. The methodology, however, could also be used in other GHG emission calculators, to improve their functionality on representing perennials.

1.1. Model definition

The Perennial-GHG model we present in this study estimates values of GHG emissions derived from the plant biomass for the entire cultivated crop area. It is a generic model that describes biomass accumulation and release, and calculates associated GHG emissions and removals. The model includes the total plant biomass: the above ground (trunk, branches, leaves and fruits) and below grown (the root system and rhizome). The model allows farm level management to be taken into account, and the system boundary is the farm gate (Hillier et al., 2011). GHG emissions arising from supplementary management options, machinery, farm electricity and goods transport need to be considered in the overall farm emissions, and for these we used the equations presented in Hillier et al., (2011) (not presented here). Regarding the below ground compartment, the model estimates plant biomass input to



Fig. 1. Model structure diagram. The emissions in plane black are positive emissions, GHGs released to the atmosphere. Emissions in grey are neutral emissions, the uptaken CO_2 equals the released CO_2. Emissions in bolt are negative emissions, atmospheric carbon fixed in the system.

the soil and subsequently decomposition. Perennial-GHG is a biomass model and does not include a soil module (which is the subject of ongoing work), so does not estimate changes in soil organic carbon (SOC). Yet, the outputs of our model can be used as inputs for a SOC model such us RothC (Coleman and Jenkinson, 1996), ECOSSE (Smith et al., 2010b), or YASSO (Liski et al., 2005).

In the Perennial-GHG model, biomass accumulation is described using different generic allometric curves, which have to be parametrized for each crop, and estimates biomass as a function of time (in years). In farmlands, most of the biomass released is due to human management interventions, such as grapping or pruning. The model specifies the contribution of each different plant part and/or residue to GHG emissions and details the annual GHG emission values. This allows investigation of the inter-annual variation in terms of biomass increment/decrease and GHGs and the contribution of each separate plant part or residue type to GHG emissions. We did not consider it necessary to take into account the effect of seasonal and inter-annual variability of climate for the following reasons: for the IBM, crop rotations are longer than 5–10 years, so positive and negative effects of the climate variability will largely cancel out over time (Harris et al., 2014). In the ABM this effect is directly accounted for by the input values of yield given by the user.

In the Perennial-GHG model, both the IBM and the ABM sub-models are comprised of different modules, which we present in the following subsections. The required model inputs are listed in Supplementary information S3. The model calculates emissions of the different GHG gases: CO_2, N_2O and CH_4. As is common-practise, the emissions from all those GHG gases are transformed into CO_2 equivalents using Global Warming Potential (GWP) values as follows:

$$CO_2eq\ (CH_4) = CH_4{*}GWP_{CH_4} \quad (1)$$

$$CO_2eq\ (N_2O) = N_2O{*}GWP_{N_2O} \quad (2)$$

$$CO_2eq = CO_2 + CO_2eq\ (CH_4) + CO_2eq\ (N_2O) \quad (3)$$

The model includes two different set of values for GWP, the widely used 2001 IPCC values (IPCC, 2001), and the most recent IPCC GWP over a 100-year time horizon presented in Myhre et al. (2013). Different values could be also specified by the user.

Information about annual GHG balance of each plant part, and for each residue, is stored in a matrix in the model. In addition, it should be noted that in the following, biomass always refers to the dry biomass, the weight of the plant excluding the water content. The percentage of C in the different plant organs is also required for the sub-models. Although not a focus of this study it should be noted that the model additional calculates the N balance in the plant.

$$Biomass = fresh\ weight{*}\ dry\ matter \quad (4)$$

where $dry\ matter = 1 - water\ content$, as a fraction of one.

$$Carbon_{organ} = Biomass_{organ}{*}Carbon\ content_{organ} \quad (5)$$

$$Nitrogen_{organ} = Biomass_{organ}{*}Nitrogen\ content_{organ} \quad (6)$$

Specific values of water, C and N content in different plant organs and species and are presented in Tables 1 and 2.

A first set of modules estimate biomass accumulation as a function of time, in which different plant parts are modelled separately and stored as annual values. The IBM defined for the woody crops therefore consists of the following modules: biomass from woody parts, leaf biomass, below ground biomass (accounting for the coarse and fine roots separately), biomass pulp for those crops that have to be de-pulped, and biomass of the yield discarded for quality reasons. This includes the total biomass produced by the plant, including all the pre-harvest biomass. In parallel, the ABM consists of modules for: above ground and stalk biomass, leaf biomass and below ground biomass (accounting for the rhizomes and roots with turnover separately). Once again, it includes all the pre-harvest biomass. Subsequently, a second set of modules estimate GHG emissions both from the plant parts and from the residues and/or the biomass naturally released from the plant. Five kinds of residue are accounted for in the IBM: litter from the leaves,

Table 1
Crop specific parameters for the individual based model (IBM), eq (11) to eq. (21). The carbon (C) and nitrogen (N) values are at harvesting time. The references of the source data are in Supplementary information S2 and figshare archive doi https://doi.org/10.6084/m9.figshare.5712127. Those tables will be interactive and updated in the future if more data are available. New versions will have new doi.

Crop	$\alpha_{1\ AGB}$	$\beta_{1\ AGB}$	$\alpha_{2\ AGBwoody}$	$\beta_{2\ AGBwoody}$	$\alpha_{3\ BGB}$	$\beta_{3\ BGB}$	$\alpha_{4\ Leaves}$	$\beta_{4\ Leaves}$	Wood dry biomass	C wood	N wood	C leaf	N leaf	Fruit dry biomass	Pulp/seed	C fruit	N fruit
Apple	0.683	1.760	0.267	2.025	0.460	1.345	0.699	0.417	0.8	0.47	0.015	0.47	0.25	0.14	–	0.47	0.0038
Citrus	0.395	2.120	0.125	2.376	0.040	2.525	1.297	0.535	0.82	0.47	0.015	0.47	0.02	0.1	–	0.47	0.0095
Cocoa	1.250	1.344	1.135	1.307	0.589	1.113	0.165	1.073	0.8	0.47	0.020	0.47	–	–	–	–	–
Coffee	3.999	0.568	3.334	0.703	0.228	1.589	0.223	0.940	0.8	0.47	0.400	0.47	0.47	0.15	0.4	0.47	1.6
Tea	1.526	0.557	1.215	0.599	0.213	0.580	0.592	0.135	0.8	0.47	0.0041	0.69	0.03	–	0	0.69	0.028
Willow	–	–	0.158	1.611	0.158	1.611	–	–	0.8	0.49	0.275	0.5	0.015	–	–	–	–
Poplar	3.389	1.605	7.223	1.257	0.781	0.745	2.426	−0.182	0.8	0.49	0.238	0.5	0.317	–	–	–	–

Table 2
Crop specific parameters for the area based model (ABM), eq (22) to eq. (28). The carbon (C) and nitrogen (N) values are at harvesting time (maturity). The references of the source data are in Supplementary information S2 new and figshare archive doi https://doi.org/10.6084/m9.figshare.5712127. Those tables will be interactive and updated in the future if more data are available. New versions will have new doi.

Crop	Stalk:AGB	AGB:roots	BGB:AGB	Stalk water content	Root senescence ratio	C stalk	N stalk	C leaf	N leaf	C root	N root
Miscanthus	0.8	0.85	0.73	0.5	0.17	0.5	0.0016	0.457	0.0045	0.41	0.015
Sugarcane	0.826		0.32	0.71	0.17	0.443	0.012	0.4525	0.014	0.405	0.00395
Switchgrass	1	0.8	0.62	0.2	–	0.44	0.003	0.462	0.01	0.44	0.03

woody parts from pruning, trees that die and the final tree cut, the fruit discarded and fruit pulp, and fine roots that die. In the ABM, three kinds of residue are accounted for: the leaves, if it is not a commodity, total above ground biomass (AGB) of the unproductive initial(s) year(s), and roots that die. The total GHG emissions from residues can be either positive or negative and this strongly depends on the residue management, which is a model input indicated by the user.

The Perennial-GHG model incorporates different residue management options. Options for wood residues are: burning, chipping followed by spreading, or chipping followed by removal. For litter, the options are either burning or litter left on the ground. For discarded fruits and pulp the management options are either: left on the ground or removed. In either case, burning will always result in positive GHG emissions but residue incorporation into the soil will result in negative emissions. If plant parts are taken away - effectively outside the farm boundary, this is considered to be neutral consistent with our farm-gate boundary (as described in the introduction), which was fixed to limit the model scope to processes over which farmers have control. Perennial-GHG allows a mix of different management techniques for each residue source, for example, 50% of the pruning residues chipped and 50% burnt.

As a final step, outputs from the modules are summed to obtain the total field level estimation of GHG emissions. The carbon in harvested products, exported beyond the farm gate is, excluded from the accounting since it is generally considered in bioenergy, food and drink sectors to be available for combustion or consumption, and thus most likely returned to the atmosphere in the short carbon cycle. However, this is not the case if is the harvested products are used to produce bio-based products such as bioplastic or bio-based building materials; these are not accounted for in the model.

For the IBM, the field CO_{2eq} is calculated by multiplying the individual value by the number of trees of each species. For monocultures, only one species is included. For intercropping or multi-cultures, the CO_{2eq} from each species is gathered:

$$Field\ CO_{2eq\ biomass\ year} = \left(\sum_{s=1}^{s=S} Ind\ CO_{2eq\ biomass\ year} * N_S\right) * A \tag{7.1}$$

where S in the number of species, $S = 1$ in monocultures. $Ind\ CO_{2eq\ biomass\ year}$ are the individual values of CO_{2eq} containing separate information about the aforementioned plant biomass and residue for each year per species s. The modules for estimated plant and residue biomass will be detailed in the forthcoming section. N_s is the number of trees per ha of each species s. This number does not equal the number of planted trees because some trees will die during the crop life period. If gapping (replacement of dead trees) is not present, then $N = N_{planted\ trees} - N_{trees\ die}$. If gapping is present, N is equal to the number of planted trees. In both cases, the percentage of trees that die is an input to the model. The model assumes a constant mortality ratio during the period: $N_{trees\ die} = N_{planted\ trees} * \%\ trees\ die/100$. A is the total cultivated area in ha.

For the ABM, the field CO_{2eq} is calculated by multiplying the per hectare value by the total area:

$$Field\ \ CO_{2eq\ biomass\ year} = \sum_{s=1}^{s=S} Area\ CO_{2eq\ biomass\ year} * A \tag{7.2}$$

where s in the number of species, $s = 1$ in monocultures $Area\ Co2eq_{mass\ year}$ are the per-ha values of CO_{2eq} containing separate information about each species s of plant biomass or residue and year. The modules for estimated plant and residue biomass will be detailed in the forthcoming section. A is the cultivated area in ha of each species.

For farms than contain both crops that fall in the ABM and the IBM categories, the field CO_{2eq} is calculated by adding the GHG derived from those crops (eq. (7.1) and eq. (7.2)).

$$Field\ CO_{2eq\ biomass\ year} = \left(Field\ CO_{2eq\ biomass\ year}\right)_{IBM} + \left(Field\ CO_{2eq\ biomass\ year}\right)_{ABM} \tag{7}$$

The annual values are then summed to derive the overall CO_{2eq} values from each plant part or residue each year of the crop lifecycle in the entire cultivated field:

$$Field\ CO_{2eq\ biomass\ year} = \sum_{year=1}^{year=\ Years} Field\ CO_{2eq\ biomass\ year} \tag{8}$$

And the overall CO_{2eq}, regardless of plant part or residues, is:

$$Field\ CO_{2eq} = \sum Field\ CO_{2eq\ biomass} \quad (9)$$

Finally, CO_{2eq} equivalent per tonne of finished product is given by:

$$CO_{2eq}\ per\ tonne\ of\ final\ product = Field\ CO_{2eq} / \sum_{year=1}^{year=\ Years} yield_{year} \quad (10)$$

where total yield is a model input.

In this section, only the equations for CO_{2eq} are shown, but a similar approach exists for individual GHGs. All the functions provide values of CO_{2eq} in kg.

Definitions of all the parameters included in the model are detailed in Table 3. The R code for the main model including all the modules is provided Supplementary information in S1 and the figshare archive doi https://doi.org/10.6084/m9.figshare.5712109. The database of empirical values used to parametrize the model is provided Supplementary information in S2 and figshare archive doi https://doi.org/10.6084/m9.figshare.5712127. The required model inputs to run the Perennial-GHG model are provided in Supplementary information S3.

2. Plant biomass modules

2.1. Individual based sub-model (IBM) for perennial woody crops

Functions in this subsection estimate biomass accumulation as a function of time in the different plant parts. They represent cumulative amounts, in units of kg per plant.

2.1.1. Biomass in wood module

This module provides the above ground biomass of the woody parts (AGBW) as a function of time. The AGBW comprises the stem plus all the branches, including twigs. Power relationships are generally used in biomass estimation (Stephenson et al., 2014) and in this case, the power law provided the best fit to the crop-growth empirical data for different crops we have (data reproduced in Supplementary information S2). The power law was not only the best fit for single crops in most cases, but also the best single function that accommodated all crops.

$$AGBW = \left(\alpha_1 age^{\beta_1}\right) * Rw_{AGB} * Rf_{AGB} \quad (11)$$

where *age* is the age of the above-ground plant part, in years. α_1 and β_1 are specific parameters (see Table 1). The Rw_{AGB} and Rf_{AGB} account for water and nutrient limitation – i.e. the growth limiting effect of lack/excess of water, and lack of fertilizers, respectively. To date, data on robust empirical Rw_{AGB} and Rf_{AGB} values for perennial crops are rare, and thus are set to 1 in the current model.

If pruning is practiced, as is common for many perennial crops, the values of AGBW are corrected to actual AGBW (actAGBW):

$$actAGBW_{year} = (AGBW - Pruning)_{year} \quad (12)$$

where *year* is the crop life year at which the plantation starts, in years, starting in 1. The parameter *age* and *year* may be the same if the plant is planted on the farm at age 0. The model allows two kinds of inputs regarding pruning values: the values can be specified either in fresh weight of pruned residues per year or as the percentage of crown removed per year.

The cumulative values of pruned biomass are:

$$AGBpruning_{year} = \sum_{year=SPrun}^{year=Years} (Pruning)_{year} + (Pruning)_{year-1} \quad (13)$$

where *SPrun* is the year in which pruning starts. This function assumes that pruning is always executed once it starts.

2.1.2. Biomass in leaves module

Two sub-models are defined for leaves, one for deciduous species and a one for evergreens. The deciduous plants module is:

Table 3
List of variables used in the Perennial-GHG model.

Variable	Meaning	Units
CO_2eq	CO_2 equivalent	kg
Biomass	Plant biomass, dry weight	kg
AGB	Above ground biomass, dry weight	kg
BGB	Below ground biomass, dry weight	kg
$Field\ CO_{2eq}$	CO_2 equivalent emissions in the farm	kg
N	Number of trees in a plantation or orchard	–
S	Number of species in the cultivated area	–
N_s	Number of trees per ha of each species *S*	–
$Ind\ CO_{2eq}$	Individual (per plant) values of biomass	–
Years	Number of years of the crop cycle = last year of the crop cycle	–
year	Each single year of the crop cycle	–
SPrun	The year in which pruning starts.	–
age	Age of the plant above ground part	year
age_{root}	Plant root age,	year
AGBW	AGB of the woody parts	kg
$actAGBW_{year}$	AGB of the woody parts after pruning	kg
$\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3$	Specific parameters for the IBM	–
Rw_{AGB}	Parameter to account for water and nutrient limitation	–
Rf_{AGB}	Parameter to account for nutrient limitation	–
l	Average lifespan of the leaves	year
SProd	The year in which production starts	–
rsen	Root senescence ratio	–
CF_{organ}	Carbon fraction in the organ	one unit
mass	Remaining mass in the decomposition model	kg
k	Decay constant in the decomposition model	–
t	Time in the decomposition model	year

$$Annual\ Leaves\ Biomass_{dec} = \alpha_2 actAGBW^{\beta_2} \quad (14.1)$$

where α_2 and β_2 are specific parameters (Table 1). Leaf biomass is therefore a function of actAGBW. eq. (14.1) is applied annually to have the annual leaf biomass. Cumulative leaf biomass is thus given by:

$$Leaf\ Biomass_{dec} = \sum_{year=1}^{year=Years} (Annual\ Leaf\ Biomass_{dec})_{year} + (Annual\ Leaf\ Biomass_{dec})_{year-1} \quad (15.1)$$

The module for evergreen plants is mathematically similar to eq. (14.1), except that the current leaf biomass does not correspond to the annual production.

$$Annual\ Leaf\ Biomass_{ev} = \alpha_2 actAGBW^{\beta_2} \quad (14.2)$$

where α_2 and β_2 are specific parameters (Table 1).

$$Discarded\ biomass_{year} = \sum_{year=SProd}^{year=Years} \left(Harvest\ yield*\%\ discarded/100\right)_{year} + (Discarded\ biomass)_{year-1} \quad (20)$$

The cumulative value of leaf biomass in this second case is:

$$Leaf\ Biomass_{ev} = \sum_{year=1}^{year=Years} Annual\ Leaf\ Biomass_{ev} + Annual\ Leaf\ Biomass_{ev/l} \quad (15.2)$$

where l is the average lifespan of the leaves.

2.1.3. Below-ground biomass module

Below-ground biomass refers to the entire root system, including both the coarse roots and the fine roots. The module to calculate root biomass is:

$$BGB = \left(\alpha_3 age_{root}^{\beta_3}\right) * Rw_{BGB} * Rf_{BGB} \quad (16)$$

where age_{root} is the plant root age, in years. The age_{root} can be equal during the first crop rotation but they will differ after biomass removal and re-growth. α_3 and β_3 are specific parameters (Table 1). This model also includes the theoretical parameters $Rw_{BGB}*Rf_{BGB}$ to account for lack and excess of water and lack of fertilizers, not parametrized yet and set equal to 1.

For estimating the percentage of fine roots as a function of plant age, the equation proposed by Kurz et al. (1996) is used. It can be seen that the proportion of fine roots (*Prop fine roots*) decreases with age:

$$Prop\ fine\ roots_{age_{root}} = 2.73 * age_{root}^{-0.841} \quad (17)$$

$$Fine\ root_{root} = Prop\ fine\ roots_{age_{root}}/100 * BGB_i \quad (18)$$

where $Prop\ fine\ roots_{age_r}$ is the proportion of fine roots at a particular plant root age, in years.

The fine roots have a short life (Withington et al., 2006). We therefore assumed the fine roots die every year and new fine roots are produced, while the coarse roots remain (Guo et al., 2006; Withington et al., 2006). The fine roots that die will either decompose to emit short cycle CO_2 or add to the soil organic carbon pool. The decomposition rate and equations are specified in the section "calculation of GHG emissions".

2.1.4. Crop yield residue module

Crop yield is not predicted in the model. It is a model input that should be indicated by the user. However, some crop yield is discarded because it does not meet required quality standards. If this is the case, the model accounts for this crop biomass, which becomes a residue instead of a commodity. The user indicates the actual harvested crop yield biomass, but the actual plant yield is:

$$Total\ yield = harvested\ yield + \left(harvested\ yield*\%\ discarded/100\right) \quad (19)$$

where % *discarded* is the percentage of unharvested yield. Hence:

where *SProd* is the year in which production starts.

A second important residue derived from the fruit is the pulp for those crops in which de-pulping is necessary, such as for coffee. The pulp biomass is calculated as a function of the yield indicated by the user. The percentage of pulp/seed is a specific parameter (Table 1).

$$Pulp\ biomass_{year} = \sum_{year=SProd}^{year=Years} \left(yield_{year}/Perc\ seed*Perc\ pulp\right)_{year} + \left(yield/Perc\ seed*Perc\ pulp\right)_{year-1} \quad (21)$$

where *Perc seed* is the percentage in one of the seeds with respect to the entire fruit (seed plus pulp). And *Perc pulp* is the percentage in the pulp with respect to the entire fruit.

2.2. Area based sub-model (ABM) for perennial grasses biomass

In the ABM, biomass values are modelled in tonnes per ha per year and may subsequently be converted to kg for consistency with the IBM model.

2.2.1. Stalk and above ground biomass module

The AGB for perennial grasses is calculated using the yield information provided by the user. The model does not predict yield but uses the provided yield information to calculate plant biomass. The user can provide the yield as either fresh plant weight, right after harvesting the plant, or plant weight after leaving it dry on the ground, along with the moisture content at that particular time or dry biomass, the plant weight excluding the water. The yield can be either the autumn or spring harvest. In this study, we have parametrized for the autumn harvest (Table 2). Two modules are defined for estimating AGB. In either case, the model considers that the plants are annually harvested and consequently a new above-

ground part grows every year. The first module should be used for those species in which the harvested part is only the stalk and the leaves are hence residues, such as sugarcane.

The annual stalk biomass is:

$$Stalk\ biomass_{age} = Yield_{age}{*}dry\ matter \tag{22}$$

where *age* is the plant aboveground age, *dry matter* is a specific values for fresh plant, given in Table 2, if the values of yield are included in the model as a fresh weight. If the yield values are input as semi-dry weight, the *dry matter* = 1 − *moisture content*. If the yield values are input as dry weight, the yield will equal the stalk biomass, hence *dry matter* = 1.

The total stalk production is hence:

$$Stalk\ biomass = \sum_{year=1}^{year=Years} (Yield_{year}{*}dry\ matter)_{year} + (Stalk\ biomass)_{year-1} \tag{23}$$

where *year* is the crop life year at which the plantation starts, in years, starting in 1 and *N* is the last year of the crop cycle. The parameter *age* and *year* may be the same if the plant is planted on the farm at age 0.

The above ground biomass:

$$AGB_{year} = Stalk\ biomass_{year}/_{stalk:AGB} \tag{24.1}$$

where *stalk* : *AGB* is the ratio, as a fraction on one, of the stalk with respect to the total AGB, a specific value (Table 2).

The cumulative values of AGB were also calculated at the end of the crop lifecycle, as in eq. (23).

In this case (eq. (24.1)), is used to calculate AGB, since the stalk biomass (from eq. (23)) and the *stalk* : *AGB* values (Table 2) are known parameters. Importantly, the plant organ ratio parameters change not only among crops, but also for the harvesting times. The model can consider those differences by using different crops specific parameters.

The second module should be used for those species in which the harvested yield includes both the stalk and the leaves, such as switchgrass.

$$AGB_{age} = Yield_{age}{*}dry\ matter \tag{24.2}$$

The cumulative values of AGB were also calculated at the end of the crop lifecycle, as in eq. (23).

Species specific values of dry matter for fresh plants are shown in Table 2. If the yield values are input as semidry weight, *dry matter* = 1 − *moisture content*. If the yield values are input as dry weight, the yield will equal the stalk biomass, hence *dry matter* = 1. In either case, if the plant is cut but not harvested in the first year(s) of it cycle, the potential yield is treated as a residue.

2.2.2. Leaf biomass module

This module estimates the biomass of leaves, in tonnes per ha and year.

$$Leaves\ biomass_{age} = AGB_{age}{*}(1 - stalk:AGB) \tag{25}$$

The cumulative values are also calculated at the end of the crop lifecycle, as in eq. (23).

When the perennial grasses harvest is after senescence, much of the life material becomes litter and is therefore considered in this section. This actually improves the quality of the harvested biomass as it has less ash and potassium without the leaves.

2.2.3. Below-ground biomass module

The below-ground biomass of the grasses comprise not only the roots but sometimes a rhizome. The rhizome is a storage organ which grows as the plant establishes, but it remains the same size in mature established crops. What we call below-ground biomass in this study includes both the rhizome and the roots, if both organs are present in the crops. Roots are about 20% of the below-ground biomass for most bioenergy crops (Dohleman et al., 2012). Previous research shows that the below-ground biomass in agricultural perennial grasses does not change appreciably over time after establishment (Dohleman et al., 2012; Ebrahim et al., 1998), and is independent of senesced rate (Amougou et al., 2011). Consequently, this sub-model assumes that from year 1 after planting, the entire root system and the rhizome are developed, and in the subsequent years the biomass of new roots is equal to the biomass of roots that senesce. For some individuals or crop varieties rhizome development may take up to three years, but the model does the aforementioned assumption for simplicity. This below-ground biomass module is always used in this form, including for the first unproductive years, if present.

The below ground biomass is hence:

$$BGB = Biomass_{roots} + Biomass_{rizhome} \tag{26}$$

The BGB module for year 1 is:

$$BGB_1 = AGB_1{*}(AGB:BGB) \tag{27}$$

where the *AGB* : *BGB* is the specific value at harvesting age, values in Table 2.

For subsequent years:

$$BGB_{ratoon\ year} = BGB_1{*}rsen \tag{28}$$

where *rsen* is the root senescence ratio, values in Table 2.

The cumulative values were also calculated at the end of the crop lifecycle, as in eq. (23). The roots that die during the year will either decompose to emit short cycle CO_2, or add to the soil organic carbon pool. The decomposition rate and equations are specified in the following section, “calculation of GHG emissions”.

2.3. Calculation of GHG emissions

Henceforth values of CO_2, N_2O and CH_4 are subsequently converted into CO_2 equivalents using equations eqs. (1)–(3).

2.3.1. Aerial biomass

The equation to estimate annual CO_2 absorbed from the atmosphere and converted into biomass from living plant parts is:

$$CO_{2\,organ} = Biomass_{organ}{*}CF_{organ}{*}\frac{44}{12}(-1) \tag{29}$$

The plant biomass values derive from the corresponding equation in section “Plant biomass modules”. CF_{organ} is the carbon fraction in the organ (Tables 1 and 2).

Plant biomass is accumulated through time, but at the end of the crop life cycle, only the root biomass prevails. The entire AGB is either harvested, i.e. if the plant is used to produce biofuel or bio-based products, or becomes residue, i.e. if the only the fruit is used, like in top-fruit trees.

2.3.2. Below-ground parts

The Perennial-GHG model does not consider root removal once the crop cycle is completed (Hastings et al., 2017), since it is a very demanding practice and is uncommon in agriculture.

Consequently, plant roots remain underground after plant harvest and become part of the soil organic carbon. Some roots die during the production period. This dead biomass will either decompose or stay as a stable component in the soil, henceforth incorporated as part of the soil organic carbon pool (Schulze and Freibauer, 2005). The roots that decompose are neutral in terms of carbon, and the remaining biomass is a negative emission accounted for in the model. It is important to note that the Perennial-GHG estimates biomass and plant residues, and derives GHGs during the crop cycle. These root soil input materials will stay in the soil for some time, depending on the soil conditions and climate (Powlson et al., 2013). Nevertheless, subsoil or tillage operations are considered in the additional management options, and the roots removed through these operations are included.

To calculate the remaining biomass of roots that die for the IBM, we used the widely-used decay function proposed by Aber et al. (1990):

$$mass = e^{-kt} \tag{30}$$

where *mass* is the remaining mass, k is the decay constant and t is the time in years. For woody crops $k = 0.51$ (Guo et al., 2006). The remaining root biomass at year i is:

$$Remaining\ mass_{\ roots\ i} = Original\ mass_{roots} * e^{-0.51\ i} \tag{31}$$

The k parameter we provide is general and can be refined for different crops and climates when robust empirical data are available.

For the ABM, root senescence is available (Table 2).

In either case, remaining biomass decreases with time and this effect is also included in the model.

The module for estimating root GHG emissions:

$$CO_2\ BGB = \Big(BGB_{end\ period} + Remaining\ mass_{roots\ \ end\ \ period}\Big) * CF_{root} * \frac{44}{12}(-1) \tag{32}$$

BGB is derived fom eq. (16) in IBM and eqs. (26)–(28) in the ABM. CF_{root} is the carbon fraction in the root, a specific parameter (Tables 1 and 2).

AGB and BGB values are fitted independently in the model. In natural plants AGB and BGB have to be considered together to account for biomass distribution and resource allocation. This is not the case for farm plants. First, management changes the above ground part and therefore overall plant carbon allocation no longer follows the natural rule. Second, and more importantly, the common practice of harvesting the AGB part but not the BGB (i.e., bioenergy crops, SRC, cropping practices in fruit trees) creates an unbalanced plant age, with the belowground system frequently older than that above ground. To reflect these differences the model needed, in turn, a separate estimator for above and belowground biomass.

2.3.3. Wood residues that are burnt

GHG emissions from burning wood residues are estimated using the following equations, presented in Akagi et al. (2011):

$$1\ Kg\ burnt\ wood\ biomass = \Big(1.509\ Kg\ CO_2 * \%\ residual\ burnt/100\Big) - wood\ biomass\ CO_2 \tag{33}$$

$$1\ Kg\ burnt\ wood\ biomass = 0.00568\ Kg\ CH_4 * \%\ residual\ burnt/100 \tag{34}$$

$$1\ Kg\ burnt\ wood\ biomass = 0.00038\ Kg\ N_2O * \%\ residual\ burnt/100 \tag{35}$$

where *wood biomass* is derived from equations eq. (13) for pruning residues or eq. (12) for the tree at the end of the cycle and/or trees that die during the period. The *% residual burnt* is the percentage of residues that are burnt. This is an input of the model (see the explanation at the beginning of section "Model definition" for details). Short cycle CO_2 stored in plant biomass as organic carbon is not accounted here as it is taken up by the plant and returned shortly after.

2.3.4. Wood residues that are chipped

If the woody parts are chipped and spread on the soil, they either add to the soil organic carbon pool (Weedon et al., 2009) or decompose to emit CO_2, which is effectively carbon neutral. To calculate the remaining soil organic carbon, we used a decay function (eq. (30)). For wood chips, the decomposition constant $k = 0.3$ (Liski et al., 2005). Hence, at year = i the remaining mass of chips is:

$$Remaining\ mass_{chip\ i} = Original\ mass_{chip} * e^{-0.3\ i} \tag{36}$$

And the module for estimating CO_2 is:

$$CO_2\ chips_i = Remaining\ mass_{chip\ i} * CF_{wood} * \%\ wood\ spread/100 * \frac{44}{12} * (-1) \tag{37}$$

where $Remaining\ mass_{chip\ i}$ is derived from eq. (36) applied after eq. (13) for pruning residues or eq. (36) applied after eq. (12) for the tree at the end of the cycle and/or trees that die during the period. CF_{wood} is the faction of carbon in the biomass (Table 1). The *% wood spread* is the percentage of the residues that are chipped and spread (see section "Model definition"). The k parameter was developed to be used in temperate climates. We use it as a general value here, but it can be refined for different crops and climates when robust empirical data are available.

If the woody parts are chipped and the chips are removed, they are regarded as neutral in terms of carbon and therefore the plant emissions are equated to zero in the Perennial-GHG model.

2.3.5. Litter burning

GHGs from litter burning are estimated using the IPCC values for biomass burnt with GHGs for agricultural residues, Table 2.5 in Chapter 2, Volume 4 of the original document (IPCC, 2006).

$$1\ Kg\ burnt\ litter\ biomass = \Big(1.515\ Kg\ CO_2 * \%\ litter\ burnt/100\Big) - wood\ biomass\ CO_2 \tag{38}$$

$$1\ Kg\ burnt\ litter\ biomass = 0.027\ Kg\ CH_4 * \%\ litter\ burnt/100 \tag{39}$$

$$1\ Kg\ burnt\ litter\ biomass = 0.00007\ Kg\ N_2O * \%\ litter\ burnt/100 \tag{40}$$

where *litter biomass* is derived in the IBM from eq. (15.1), in the case

of deciduous species and eq. (15.2) for evergreen species. *litter biomass* is derived in the ABM from eq. (25) for litter or eq. (24) for the unproductive year. From the combustion, CO_2, N_2O and CH_4 are produced. Values of those gases are transformed into CO_2eq using equations eqs. (1)–(3). The *% litter burnt* is the percentage of residues that go to the burnt set (see section "Model definition").

2.3.6. Litter left on the ground

When the leaves are left on the ground, they either decompose or become part of the soil organic carbon pool (Schulze and Freibauer, 2005). The litter that decomposes is carbon neutral. To calculate the remaining soil organic carbon we used the decay function eq. (23). In the IBM, the decomposition value for litter $k = 0.83$ (Wu et al., 2012). In the ABM, the decomposition value $k = 0.776$ (Amougou et al., 2012).

The equation to estimate CO_2 from litter is:

$$CO_2\ litter_i = Remaining\ mass_{litter\ i} * CF_{leaves} * \frac{44}{12} * (-1) * \%\ litter\ left / 100 \quad (41)$$

where $Remaining\ mass_{litter\ i}$ is the mass after using eq. (15) for calculating litter biomass followed by eq. (22) for calculating litter decomposition in the IBM sub-model and eq. (25) for litter biomass followed by eq. (23) for litter decomposition in the ABM sub-model. CF_{leaves} is the carbon fraction in the leaves, a specific value (Tables 1 and 2). The *% litter left* is the proportion of litter left on the ground (see section "Model definition").

2.3.7. Discarded fruits left on the ground

Some produce which does not meet quality standards may be left on the ground instead of harvested. If this is the case, it either decomposes or becomes part of the soil organic carbon pool. The part that decomposes is carbon neutral. To calculate the remaining soil organic carbon we used the decay function eq. (30). The fruit decomposition value $k = 0.83$ (Wu et al., 2012).

The equation to estimate CO_2 from those fruits is:

$$CO_2\ fruits = Remaining\ mass_{discarded\ fruit} * CF_{fruit} * \%\ fruit\ dis / 100 * \frac{44}{12} * (-1) \quad (42)$$

The biomass of discarded fruits is calculated using eq. (20). CF_{fruit} is the carbon fraction in the fruits, a specific value (Table 1). The *% fruit disc* is the percentage of discarded fruits, a model input.

2.3.8. Fruit pulp left on the ground

If the pulp of de-pulped fruits is spread out on the farm, it either decomposes or becomes part of the soil organic carbon pool. The part that decomposes is carbon neutral. To calculate the remaining soil organic carbon we used the decay function eq. (23). The fruit decomposition value $k = 0.83$.

The equation to estimate CO_2 from those fruits is:

$$CO_2\ fruits = Remaining\ mass_{pult} * CF_{fruit} * \%\ pulp / 100 * \frac{44}{12} * (-1) \quad (43)$$

The biomass of discarded fruits is calculated using eq. (21). CF_{fruit} is the carbon fraction in the fruits, a specific value (Table 1). The *% pulp* is the percentage of pulp that is spread out, a model input.

2.3.9. Composting residues from leaves, wood chips, discarded fruits and pulp

If the residues are composted within the farm, to be used either in the farm or in a different area, the model accounts for the GHGs. If the residues are removed for composting elsewhere, then they are considered GHG neutral. Although plant residues accumulate biomass, GHGs are emitted during composting. Those GHGs result from fuel used in combustion and from the degradation of the feedstock biomass (Boldrin et al., 2009; Brown et al., 2008). GHGs from the fuel from combustion and the degradation depend on the type of technology used in composting (Brown et al., 2008). The equation to estimate CO_2 from composting is:

$$CO_2eq\ compost = CO_{2eq}\ Biomass_{compost} + CO_{2eq}\ Compost_{process} + CO_{2eq}\ Compost_{energy} \quad (44)$$

The $CO_2\ Biomass_{compost}$ can be calculated

$$CO_2\ Biomass_{compost} = \left(biomass_{residue} * CF_{residue} * \frac{44}{12} * \left(1 - \%C_{degraded} / 100 \right) \right) \quad (45)$$

where $\%C_{degraded}$ is the percentage of carbon that degrades during the process of decomposition. The model uses the values of $\%C_{degraded} = 60$ for open systems and $\%C_{degraded} = 55$ for enclosed systems (Boldrin et al., 2009).

To estimate the $CO_{2eq}\ Compost_{process}$, the model uses the mean value of the range of compost emission factors presented in Boldrin et al. (2009) and the values to calculate CO_2eq from CH_4 and N_2O from eqs. (1)–(3). The compost emissions factor vary between open and enclosed technology:

$$CO_2eq\ (CO_2)open = biomass_{residue} * (1 + WC_{residue}) * 0.25 \quad (46)$$

$$CO_2eq\ (CO_2)enclosed = biomass_{residue} * (1 + WC_{residue}) * 0.3 \quad (47)$$

$$CO_2eq\ (CH_4)open = biomass_{residue} * (1 + WC_{residue}) * 0.0035 * 34 \quad (48)$$

$$CO_2eq\ (CH_4)enclosed = biomass_{residue} * (1 + WC_{residue}) * 0.0009 * 34 \quad (49)$$

$$CO_2eq\ (N_2O)open = biomass_{residue} * (1 + WC_{residue}) * 0.001 * 298 \quad (50)$$

$$CO_2eq\ (N_2O)enclosed = biomass_{residue} * (1 + WC_{residue}) * 0.00659 * 298 \quad (51)$$

where $WC_{residue}$ is the fraction of water in the introduced residue. It was necessary to consider the water since the emission factors were based on feedstock wet weight.

To estimate the $CO_{2eq}\ Compost_{energy}$, the model used the diesel intake consumption factor presented in Boldrin et al. (2009), which is approximately 3 L per kg of wet residue for both open and enclosed technology. The emission factor for combustion of diesel is 2.7 kg CO_2eq/litre (Fruergaard et al., 2009). Therefore:

$$CO_{2eq}\ Compost_{energy} = biomass_{residue} * (1 + WC_{residue}) * 8.1 \quad (52)$$

2.4. Model parametrization

The generic model needs empirical data for parametrization to be functional and applicable for different crops, different varieties, and different geographic regions. The required empirical data for

parameterization are biomass quantity of the different plant parts at different age. The most accurate method to obtain plant biomass values is by destructive sampling (see Chave et al., 2015), but if these are not available, local allometric equations to estimate biomass as a function of plant size can be used, for example the ratio of height to biomass in Miscanthus (Kalinina et al., 2017).

Empirical values of biomass of the different plant parts at different ages are then fitted to a power law equation. We used the nonlinear least-squares estimates for parameter estimation, using the R build in function "nls" (R code in Supplementary information S1). The generic model needs empirical data not only to work for most crops, but also to improve the current estimates presented in Tables 1 and 2, and to account for varietal and geographical differences. The data used for parametrize the crops is in Supplementary information S2.

The power law is frequently used for biomass estimation of woody plants (Stephenson et al., 2014). This function is asymptotic for small alpha values, as in the present case (Table 2). In addition, tree biomass in the model is highly related to the management practices which reduce biomass (i.e. pruning), and therefore unlimited growth.

2.5. Case studies: biomass and GHGs in four main crops: apple, coffee, Miscanthus, and sugarcane

The perennial-GHG model presented in section 2 is used here to estimate GHGs in four perennial systems: apple, coffee, *Miscanthus* and sugarcane. We selected these crops to have a variety of temperate, tropical, food and bioenergy examples. In each case, we calculated GHGs in a standard 1 ha production area. We used the Myhre et al. (2013) GWP over a 100-year time horizon. We then used the Cool Farm Tool (Hillier et al., 2011) to calculate GHGs due to agrochemicals, fertilizers and energy consumed during crop management for those example using representative management practices. Our aim here is to illustrate the model application using typical management practices (Table 4), and also to examine the importance of the biomass pool in the context of total GHG emissions from crop production. We used specified values at crop maturity. In every case, further transportation of the crop was excluded from this analysis, consistent with our farm gate boundary.

The negative GHG emissions derived from the plant biomass exceed the positive GHG emissions from the supply of nutrients and agrochemicals, resulting in negative overall emissions (Fig. 2). In coffee and sugarcane the total emissions are positive due to the litter and final cut burning. For the perennial grasses, sugarcane and *Miscanthus*, most of the negative GHGs are due to root biomass accumulation followed by litter left on the ground. The amount of litter is larger but it mainly decomposes in the following years (Schulze and Freibauer, 2005) while the root biomass persists for longer. In the top-fruit crops, apple and coffee, most of the negative GHGs are due to root biomass accumulation. Litter and residues left on the ground also contribute to sink carbon in the top-fruit crops, but to a lesser extent. Litter is less abundant and decomposes faster than for the bioenergy crops. For sugar cane especially, emissions are substantial during the crop lifecycle, mainly as a result of residue burning. If burning is avoided in sugarcane and coffee, these crops would have had large negative values, in spite of the fact that these crops require more nutrient supply than the others. This illustrates that alternative practices may significantly impact GHG emissions. A large source of negative GHGs could have been obtained from sugarcane, coffee and apple with different management. Nevertheless, in every case, the results show that leaving the roots and the removed leaves on the ground contributes to fixing atmospheric carbon,

Table 4
Farm and crop parameters used in the case examples.

Crop	Production tonnes per ha[a]	Lifespan years	N trees per ha	Residue Management[a]					Fertilizers kg per ha[a]			Agrochemicals		Energy consumed annually
				First years discarded	Litter	Pruning	Discarded fruits	Fruit pulp	Trees end cycle	Nitrogen	Potassium	Phosphorus	Pesticides	Herbicides	
Apple	200 wet	20	800	–	100% left on the ground	chipped, 20% left on the ground and 80% removed	left on the ground	–	cut and removed	67 annually	70 every two years	90 every two years	Annually applied	–	2000 MJ
Coffee	2.5 wet	20	1500	–	100% left on the ground	chipped, 20% left on the ground and 80% removed	20% left on the ground and 80% composted. Compost taken away	100% composted. Compost taken away	cut and burnt	300 annually	50 annually	25 annually	Annually applied	–	1000 MJ
Miscanthus	25–40 (20% hum)	15	–	100% left on the ground	100% left on the ground	–	–	–	–	–	–	–	–	Applied Year 1	1050 MJ
Sugarcane	70–120	6	–	100% left on the ground	80% burnt and 20% left on the ground	–	–	–	–	70 annually	60 annually	90 annually	Annually applied	Applied Year 1	1500 MJ

[a] Production, residues and fertilizers vary among years. The values presented in this table are values are at crop maturity.

providing noticeable negative GHGs. Interestingly; the C input in the soil at the end of crop cycle was 8–10 tonnes for all crops. It is important to mention that the root and litter biomass input in the soil is not equivalent to the carbon sink in the soil. The quantity of carbon that stays in the soil depends not only on the input, put also on the former land use and soil properties (Dixon et al., 1994; Don et al., 2011). Evaluating such soil processes is beyond the scope of this study and it requires the use of process based models of soil biochemistry.

The annual contribution of each plant residue and fertilizer can be seen in Fig. 3 for the case of apple and *Miscanthus*. In apple, plant biomass and residue carbon accumulation increase exponentially with time (Fig. 3, left). Most of the negative GHGs are due to biomass accumulation in the woody part of the tree. But those potential negative emissions become neutral when the trees are removed. Chips and litter also contribute to the fixation of some atmospheric carbon, but a large proportion of their biomass may decompose in the future. However, GHGs from chips have a longer life and contain more carbon and stable compounds than litter, contributing to longer term carbon storage. That characteristic produces a carbon accumulation curve with a marked decreasing slope. The GHG emissions due to fertilizers applied every 2 years are fairly constant through the life of the crop. Our model estimates a total negative value of −360 $MgCha^{-1}$, stored after 20 years, similar to the range value of −230 to −475 $MgCha^{-1}$ after 20 years measured by Wu et al. (2012). The root biomass and the aerial woody biomass measured in that study were 22.93 Mg ha^{-1} and 125 Mg ha^{-1}, respectively, while the root and aerial woody biomass predicted in our model were 25.4 Mg ha^{-1} and 105 respectively.

In *Miscanthus*, the first year growth material left on the ground - including both the leaves and the stalk - is almost totally decomposed in 8 years (Fig. 3, right). Plant residues left on the ground from other years also contribute to the carbon pool, but we expect that they decompose in about 8 years, as the residues of the first year did. Hence, they may not have a very long term impact in terms of carbon, but still they have a slight contribution to negative GHGs in the long term. This rapid biomass loss causes a decrease in the cumulative litter curve (Fig. 3, right). The annual biomass litter production of 5–7.6 Mg ha^{-1} $year^{-1}$ derived from our model is the same annual value of 5–7.5 Mg ha^{-1} measured form field in Robertson et al. (2016). The annual soil organic carbon inputs from the roots was 2.12 Mg ha^{-1} $year^{-1}$, similar to the value of 2–3 Mg ha^{-1} $year^{-1}$ showed in Dondini et al. (2009), Zatta et al. (2012), and Zimmerman et al. (2013). Once again, our model provided similar values to those measured in the field, confirming the suitability of the model for both perennial bioenergy and food crops.



Fig. 2. CO_{2eq} emission in Mg at the end of the crop cycle per plant organ, residue and agrochemical for (a) an apple orchard, (b) a coffee plantation, (c) a *Miscanthus* field and (d) a sugarcane field. Details of farm management are detailed in Table 4.



Fig. 3. Annual CO_{2eq} emissions in Mg at the end of the crop cycle per plant organ, residue and agrochemical in an apple orchard with a life period of 20 years. Details of farm management are detailed in Table 4.

3. Discussion

Quantifying CO_2 capture by plants and biomass accumulation and changes in soil carbon, are key in evaluating the impacts of perennial crops in life cycle assessment. We have presented the Perennial-GHG, a working model that can be used to assess the contribution of biomass to GHGs in perennial crops. It is applicable both to food and bioenergy crops, and we have already parameterized it for several crops (Tables 1 and 2). We used the model to calculate GHGs in four perennial systems as an illustration. In every case, the carbon stored in plants due to biomass accumulation and derived plant residues more than offsets the contribution of agrochemicals and nutrients (Fig. 2). This finding is timely, and highlights the importance of taking into consideration crop biomass of perennial plants as contributors to climate change mitigation. This model will help to reduce the uncertainty that exists in quantifying the benefits of perennial crops. In addition, the model supports the FAO's drive toward "perennialization" or increase of perennial crops strategy (Rai et al., 2011), to help to mitigate climate change and increase food and ecosystem security (Glover et al., 2010).

The Perennial-GHG is a theoretical model that needs empirical data to be parametrized. Henceforth, most of the uncertainty and errors are linked with the variability of the empirical data and not with the model definition itself. Therefore, model uncertainty and sensitivity cannot be quantified in this paper because it depends on the existing empirical data. Most of our data sources did not show standard deviation of the empirical measurements, either for the biomass or decomposition values. For that reason, uncertainly was not specified and accounted for in this paper. Adding more empirical data and re-defining the parameters in a more precise way may improve the model and reduce uncertainty. Indeed, the Perennial-GHG model can be parametrized at farm level but this will require within-farm experiments and biomass measurements, which will incur additional costs. Additionally, it is important to bear in mind that GHGs from other overlooked sources, i.e. harvesting operations, machinery emissions, commodity transportation and storage or GHGs derived from plant reproduction, have been excluded in this analyses. To derive the total crop GHG balance, they should also be accounted for. As yield is not estimated in the model, for theoretical or research purposes crop-production models can be used to estimate yield, which can be then used as an input in the presented model. Examples of such models are the Miscanfor model for *Miscanthus* (Hastings et al., 2009) or the Yield-SAFE model for tree crops (van der Werf et al., 2007).

The presented Perennial-GHG model could be improved in several ways in the future which we could not consider here due to the lack of empirical data. First, geographic or climate differences among and within crops have not been considered in the proposed model, despite acknowledgement that climate can affect both plant growth and residue decomposition (Basso et al., 2017). Regarding plant growth, we used published empirical data to parametrize the model from the current area of distribution of the considered crop (reproduced in Supplementary information S2). We aim to model crops inside their potential distribution area, and hence discard unlikely production scenarios. Disregarding the effect of climate on decomposition rate is a more important consideration. Nonetheless, for wood decomposition, the effect of climate is a secondary factor (Bradford et al., 2014), and litter has a short decomposition period regardless of location (Schulze and Freibauer, 2005). In any case, the Perennial-GHG model allows different regional

decomposition parameters, although we did not explore those in this study. In a similar way, the Perennial-GHG model has a combustion parameter for woody residues (eq. (31)–(33)) and the IPCC model for combustion parameters for agricultural residues (eq. (36)–(38)), which is used for litter and bioenergy crop burning. Those parameters could be refined in the future, if more empirical data is acquired. Similarly, GHG emissions from composting can be refined in the future as the model considers only main basic technologies (Boldrin et al., 2009). The effect of lack or excess of fertilizer and water was included as a parameter in the IBM model but it was not parameterized due to the lack of robust empirical data (see section 2.1.1 for more details). Different mortality ratios among climates are already considered in the model: in the IBM mortality is a model input; in the ABM mortality is a directly reflected in the yield, a model input. Seasonal variations in terms of plant growth and residue production also exist. However, it was not necessary to include them in the IBM model since the model evaluates annual and not seasonal biomass, residues and GHGs. For the AMB, the biomass ratios change among seasons (Amougou et al., 2012). This is currently considered by requiring as input the harvest period in the model (Table 2). Besides, no varietal differences within crops have yet been considered. We pooled the data of different varieties for each crop, due to the lack of robust data of different varieties. Once again, the present model allows future inclusion of different parameters for different varieties. Once robust data exist, that information can and should be incorporated into the model.

The Perennial-GHG presented in this paper estimates the plant carbon output during the crop cycle, since the plant is established in the ground until it is harvested, and not beyond. It is important to bear in mind that the model does not estimate the persistence of carbon after it leaves the farm gate (see details in the model definition section). At the final harvest, some litter and roots are still in the ground in organic forms and over time will decompose, releasing a fraction of the stored C. Litter and fine roots have, in general, a short life span, thus the C released will occur in the following years. On the other hand, woody roots are quite stable and will decompose slowly (Guo et al., 2006; Withington et al., 2006). The carbon finally stored will depend on the soil and environmental conditions (Dondini et al., 2009) and subsequent land use. The stability of the carbon in the system is highly dependent on the existing carbon in the system, and on the land use after the perennial cultivation. The capacity to store carbon, and it's persistence in the soil, depends on the soil C concentration before the plantation, and on the climate (Powlson et al., 2013). The model also calculates the nitrogen accumulated in the different organs in the plant. This is not required for estimating GHGs, but it gives information about the nitrogen cycle that may be useful for other purposes, such as in studies of nutrient balance. A soil organic carbon model is currently being implemented alongside this biomass model. Both together are required to estimate GHGs and carbon balance from perennial crops. These models will be incorporated in to the Cool Farm Tool (Hillier et al., 2011).

Data and software availability

The R code for the main model including all the modules is provided in Supplementary information S1 and the figshare archive doi https://doi.org/10.6084/m9.figshare.5712109 (https://figshare.com/articles/Perennial-GHG_code_/5712109/1). The database of empirical values used to parametrize the model is provided in Supplementary information S2 and figshare archive doi https://doi.org/10.6084/m9.figshare.5712127 (https://figshare.com/articles/Data_-_crop_biomass_as_a_funcion_of_crop_age_-_empirical_data_for_perennial_crops/5712127). The required model inputs to run the Perennial-GHG model are provided in Supplementary information S3.

Funding

This work was funded by NERC under the project (NE/N017854/1).

Acknowledgements

We gratefully acknowledge the support of the Cool Farm Alliance and Terravesta Ltd in co-designing the model. Thanks to the anonymous reviewers and Editor for improving the first version of this manuscript.

Appendix A. Supplementary data

Supplementary data related to this article (S1, S2 and S3) can be found at https://doi.org/10.1016/j.envsoft.2017.12.005.

References

Aber, J.D., Melillo, J.M., McClaugherty, C.A., 1990. Predicting long-term patterns of mass loss, nitrogen dynamics, and soil organic matter formation from initial fine litter chemistry in temperate forest ecosystems. Can. J. Bot. 68, 2201–2208.
Akagi, S., Yokelson, R.J., Wiedinmyer, C., et al., 2011. Emission factors for open and domestic biomass burning for use in atmospheric models. Atmos. Chem. Phys. 11, 4039–4072.
Amougou, N., Bertrand, I., Machet, J.M., Recous, S., 2011. Quality and decomposition in soil of rhizome, root and senescent leaf from Miscanthus x giganteus, as affected by harvest date and N fertilization. Plant Soil 338, 83–97.
Amougou, N., Bertrand, I., Cadoux, S., Recous, S., 2012. Miscanthus × giganteus leaf senescence, decomposition and C and N inputs to soil. GCB Bioenergy 4, 698–707.
Basso, B., Dumont, B., Maestrini, B., Shcherbak, I., Robertson, G.P., Porter, J.R., Smith, P., et al., 2017. Reduced crop residues returns and rising temperatures induce soil carbon loss, exacerbating yield declines. Nat. Clim. Change.
Boldrin, A., Andersen, J.K., Møller, J., Christensen, T.H., Favoino, E., 2009. Composting and compost utilization: accounting of greenhouse gases and global warming contributions. Waste Manag. Res. 27, 800–812.
Bradford, M.A., Warren II, R.J., Baldrian, P., et al., 2014. Climate fails to predict wood decomposition at regional scales. Nat. Clim. Change 4, 625–630.
Brandão, M., Levasseur, A., Kirschbaum, M.U., et al., 2013. Key issues and options in accounting for carbon sequestration and temporary storage in life cycle assessment and carbon footprinting. Int. J. Life Cycle Assess. 18, 230–240.
Brown, S., Kruger, C., Subler, S., 2008. Greenhouse gas balance for composting operations. J. Environ. Qual. 37, 1396–1410.
Chave, J., Réjou-Méchain, M., Búrquez, A., Chidumayo, E., Colgan, M.S., Delitti, W.B., Duque, A., Eid, T., Fearnside, P.M., Goodman, R.C., Henry, M., 2014. Improved allometric models to estimate the aboveground biomass of tropical trees. Global change biology 20, 3177–3190.
Coleman, K., Jenkinson, D.S., 1996. RothC-26.3-A Model for the turnover of carbon in soil. In: Powlson, D.S., Smith, P., Smith, J.U. (Eds.), Evaluation of Soil Organic Matter Models. Springer, Berlin, Heidelberg, pp. 237–246.
Cox, T.S., Glover, J.D., Van Tassel, D.L., Cox, C.M., DeHaan, L.R., 2006. Prospects for developing perennial grain crops. Bioscience 56, 649–659.
Dawson, J.J.C., Smith, P., 2007. Carbon losses from soil and its consequences for land management. Sci. Total Environ. 382, 165–190.
Davis, S.C., Parton, W.J., Del Grosso, S.J., Keough, C., Marx, E., Adler, P.R., DeLucia, E.H., 2012. Impact of second-generation biofuel agriculture on greenhouse-gas emissions in the corn-growing regions of the US. Front. Ecol. Environ. 10, 69–74.
Dixon, R., Brown, S., Houghton, R.E.A., Solomon, A.M., Trexler, M.C., Wisniewski, J., 1994. Carbon pools and flux of global forest ecosystems. Science 263, 185–189.
Dohleman, F.G., Heaton, E.A., Arundale, R.A., Long, S.P., 2012. Seasonal dynamics of above-and below-ground biomass and nitrogen partitioning in Miscanthus × giganteus and Panicum virgatum across three growing seasons. GCB Bioenergy 4, 534–544.
Dohleman, F.G., Long, S.P., 2009. More productive than maize in the Midwest: how does Miscanthus do it? Plant Physiol. 150, 2104–2115.
Don, A., Schumacher, J., Freibauer, A., 2011. Impact of tropical land-use change on soil organic carbon stocks–a meta-analysis. Global Change Biol. 17, 1658–1670.
Dondini, M., Hastings, A., Saiz, G., Jones, M.B., Smith, P., 2009. The potential of Miscanthus to sequester carbon in soils: comparing field measurements in Carlow, Ireland to model predictions. GCB Bioenergy 1, 413–425.
Ebrahim, M.K., Zingsheim, O., El-Shourbagy, M.N., Moore, P.H., Komor, E., 1998. Growth and sugar storage in sugarcane grown at temperatures below and above the optimum. J. Plant Physiol. 153, 593–602.
Fruergaard, T., Astrup, T., Ekvall, T., 2009. Energy use and recovery in waste management and implications for accounting of greenhouse gases and global warming contributions. Waste Manag. Res. 27, 724–737.
Glover, J.D., Reganold, J., Bell, L., et al., 2010. Increased food and ecosystem security

via perennial grains. Science 328, 1638–1639.
Guo, L., Halliday, M., Gifford, R., 2006. Fine root decomposition under grass and pine seedlings in controlled environmental conditions. Appl. Soil Ecol. 33, 22–29.
Harris, I.P., Jones, P.D., Osborn, T.J., Lister, D.H., 2014. Updated high-resolution grids of monthly climatic observations–the CRU TS3. 10 Dataset. Int. J. Climatol. 34, 623–642.
Hastings, A., Clifton-Brown, J., Wattenbach, M., Mitchell, C.P., Smith, P., 2009. The development of MISCANFOR, a new Miscanthus crop growth model: towards more robust yield predictions under different climatic and soil conditions. GCB Bioenergy 1, 154–170.
Hastings, A., Mos, M., Yesufu, J.A., McCalmont, J., Shafei, R., Ashman, C., Nunn, C., Scheule, H., Cosentino, S., Scalici, G., Scordia, D., Wanger, M., Clifton-Brown, J., 2017. Economic and environmental assessment or seed and rhizome propagated *Miscanthus* in the UK. Front. Plant Sci. 8 https://doi.org/10.3389/fpls.2017.01058.
Hillier, J., Walter, C., Malin, D., Garcia-Suarez, T., Mila-i-Canals, L., Smith, P., 2011. A farm-focused calculator for emissions from crop and livestock production. Environ. Model. Software 26, 1070–1078.
IPCC, 2001. Climate Change 2001: the Scientific Basis. Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK.
IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change.
Kalinina, O., Nunn, C., van der Weijde, T., Ozguven, M., Tarakanov, I., Schüle, H., Trindade, L.M., et al., 2017. Extending *Miscanthus* cultivation with novel germplasm at six contrasting sites. Front. Plant Sci. 8. doi.org/10.3389/fpls.2017.00563.
Kurz, W.A., Beukema, S.J., Apps, M.J., 1996. Estimation of root biomass and dynamics for the carbon budget model of the Canadian forest sector. Can. J. For. Res. 26, 1973–1979.
Liski, J., Palosuo, T., Peltoniemi, M., Sievänen, R., 2005. Carbon and decomposition model Yasso for forest soils. Ecol. Model. 189, 168–182.
McCalmont, J.P., Hastings, A., McNamara, N.P., Richter, G.M., Robson, P., Donnison, I.S., Clifton-Brown, J., 2015. Environmental costs and benefits of growing Miscanthus for bioenergy in the UK. GCB Bioenergy 9, 489–507.
Myhre, G., Shindell, D., Bréon, F.M., Collins, W., Fuglestvedt, J., Huang, J., Koch, D., Lamarque, J.F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., Zhang, H., 2013. Anthropogenic and natural radiative forcing. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), Climate Change 2013: the Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
Paustian, K., Six, J., Elliott, E.T., Hunt, H.W., 2000. Management options for reducing CO2 emissions from agricultural soils. Biogeochemistry 48, 147–163.
Powlson, D.S., Smith, P., De Nobili, M., 2013. Soil organic matter. Chapter 4. In: Gregory, P.J., Nortcliff, S. (Eds.), Soil Conditions and Plant Growth. Blackwell, Oxford, pp. 86–131.
Priesack, E., Gayler, S., 2009. Agricultural crop models: concepts of resource acquisition and assimilate partitioning. In: Progress in Botany. Springer, pp. 195–222.
Rai, M., Reeves, T., Pandey, S., Collette, L., 2011. Save and Grow: a Policymaker's Guide to Sustainable Intensification of Smallholder Crop Production. FAO, Rome.
Robertson, G.P., Paul, E.A., Harwood, R.R., 2000. Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. Science 289, 1922–1925.
Robertson, A.D., Whitaker, J., Morrison, R., Davies, C.A., Smith, P., McNamara, N.P., 2016. A Miscanthus plantation can be carbon neutral without increasing soil carbon stocks. GCB Bioenergy. https://doi.org/10.1111/gcbb.12397.
Schulze, E.D., Freibauer, A., 2005. Environmental science: carbon unlocked from soils. Nature 437, 205–206.
Smith, J.U., Gottschalk, P., Bellarby, J., Chapman, S., Lilly, A., Towers, W., Bell, J., Coleman, K., Nayak, D.R., Richards, M.I., Hillier, J., 2010a. Estimating changes in national soil carbon stocks using ECOSSE-a new model that includes upland organic soils. Part I. Model description and uncertainty in national scale simulations of Scotland. Clim. Res. 45, 179–192.
Smith, P., Martino, D., Cai, Z., et al., 2008. Greenhouse gas mitigation in agriculture. Phil. Trans. Biol. Sci. 363, 789–813.
Smith, P., Gregory, P.J., Van Vuuren, D., et al., 2010b. Competition for land. Phil. Trans. Biol. Sci. 365, 2941–2957.
Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E.A., Haberl, H., Harper, R., House, J., Jafari, M., Masera, O., Mbow, C., Ravindranath, N.H., Rice, C.W., Robledo Abad, C., Romanovskaya, A., Sperling, F., Tubiello, F., 2014. Agriculture, forestry and other land use (AFOLU). In: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner, S., Eickemeier, P., Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., Zwickel, T., Minx, J. (Eds.), Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 811–922.
Stephenson, N.L., Das, A., Condit, R., et al., 2014. Rate of tree carbon accumulation increases continuously with tree size. Nature 507, 90–93.
Tubiello, F.N., Salvatore, M., Rossi, S., Ferrara, A., Fitton, N., Smith, P., 2013. The FAOSTAT database of greenhouse gas emissions from agriculture. Environ. Res. Lett. 8, 015009.
van der Werf, W., Keesman, K., Burgess, P., Graves, A., Pilbeam, D., Incoll, L.D., Metselaar, K., Mayus, M., Stappers, R., van Keulen, H., Palma, J., 2007. YieldSAFE: a parameter-sparse, process-based dynamic model for predicting resource capture, growth, and production in agroforestry systems. Ecol. Eng. 29, 419–433.
Weedon, J.T., Cornwell, W.K., Cornelissen, J.H., Zanne, A.E., Wirth, C., Coomes, D.A., 2009. Global meta-analysis of wood decomposition rates: a role for trait variation among tree species? Ecol. Lett. 12, 45–56.
Whitaker, J., Ludley, K.E., Rowe, R., Taylor, G., Howard, D.C., 2010. Sources of variability in greenhouse gas and energy balances for biofuel production: a systematic review. GCB Bioenergy 2, 99–112.
Whittaker, C., McManus, M.C., Smith, P., 2013. A comparison of carbon accounting tools for arable crops in the United Kingdom. Environ. Model. Software 46, 228–239.
Withington, J.M., Reich, P.B., Oleksyn, J., Eissenstat, D.M., 2006. Comparisons of structure and life span in roots and leaves among temperate trees. Ecol. Monogr. 76, 381–397.
Wollenberg, E., Tapio-Bistrom, M., Grieg-Gran, M., 2013. In: Wollenberg, E., Tapio-Bistrom, M., Grieg-Gran, M., Nihart, A. (Eds.), Climate Change Mitigation and Agriculture. Routledge, pp. 3–27.
Wu, T., Wang, Y., Yu, C., Chiarawipa, R., Zhang, X., Han, Z., Wu, L., 2012. Carbon sequestration by fruit Trees-Chinese apple orchards as an example. PLos One 7 e38883.
Yan, X., Yagi, K., Akiyama, H., Akimoto, H., 2005. Statistical analysis of the major variables controlling methane emission from rice fields. Global Change Biol. 11, 1131–1141.
Zatta, A., Hastings, A., Clifton-Brown, J., Robson, P., Monti, A., 2012. Land use change from grassland to Miscanthus: effects on soil carbon content and estimated mitigation benefit after 6 years. GCB-Bioenergy 6–4, 360–370.
Zimmerman, J., Styles, D., Hastings, A., Dauber, J., Jones, M.B., 2013. Assessing the impact of within crop heterogeneity ("patchiness") in young Miscanthus x giganteus fields on economic feasibility and soil carbon sequestration. Global Change Biology Bioenergy 6–5, 566–576.